



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

by reaction in more than half the roots, and relaxation index by reaction in fewer than half. The accompanying table from the article summarizes the results:

	Presentation time	Critical time	Reaction time	Relaxation index
Lepidium sativum at 17-18C. .	5.5 min.	6 min.	25.5 min.	30
“ “ at 25-27C. .	1.5 “	2 “	12.5 “	40
Lupinus albus at 17-18C. .	8.5 “	11 “	46.5 “	20
“ “ at 25-27C. .	2.0 “	7 “	33.0 “	25

One is struck here by the time demanded for relaxation from an exposure; it is very much greater than reported by FITTING. The author attempted, without full success, to develop a formula by which any one of the critical periods can be calculated from the other three.—WILLIAM CROCKER.

Photosynthesis.—LUBIMENKO¹⁴ finds that there is a light optimum for the production of dry substance by green plants. The absolute value of this optimum is less than that by which the chlorophyll apparatus is able to furnish the maximum of photochemical work expressed in the decomposition of CO₂. By means of monochromatic filters and gasometric determinations, a comparison is made of the action of the different colored rays on the decomposition of CO₂. These results are compared with the action of the same rays on the production of total dry weight. The energy for CO₂ decomposition in colored light depends upon the absorption of the various colored rays by the chloroplasts as well as on their caloric energy. The author objects to the method employed by KNIEP and MINDER and others in determining the influence of different colored rays on CO₂ assimilation, on the ground that they measured the quantity of light falling upon the leaf and not that absorbed by the chloroplasts. The real carbon fixation expressed by the increase in dry weight is influenced unequally by the different colored rays. The maximum increase in dry weight occurs under the action of the blue-violet rays. The yellow-orange rays are inferior to the red rays, and the minimum occurs in the green rays. It is necessary to assume two successive stages in the photosynthetic process. The first is characterized by the decomposition of CO₂ and synthesis of the first organic product. In this stage of the process the plant utilizes predominantly the energy of the red rays of the solar spectrum. The second stage is characterized by the definitive fixation of the first organic product, and the plant employs for this work especially the blue-violet rays.—CHAS. O. APPLEMAN.

¹⁴ LUBIMENKO, M. W., L'assimilation chlorophyllienne et la production de substance sèche à la lumière blanche et à la lumière colorée. Rev. Gén. Bot. 23:1-14. 1911.